




# Recommended Immunization Schedule Ages Birth – 6 Years


▼ Vaccine	Age ▶	Birth	1 Month	2 Months	4 Months	6 Months	12 Months	15 Months	18 Months	19-23 Months	2 - 3 Years	4 - 6 Years
Hepatitis B		HepB	HepB			HepB						
Rotavirus				RV	RV	RV						
Diphtheria, Tetanus, Pertussis				DTaP	DTaP	DTaP	DTaP	DTaP				DTaP
<i>Haemophilus influenzae</i> type b				Hib	Hib	Hib	Hib					
Pneumococcal				PCV	PCV	PCV	PCV				PPSV	
Inactivated Poliovirus				IPV	IPV	IPV						IPV
Influenza						Influenza (Yearly)						
Measles, Mumps, Rubella							MMR					MMR
Varicella							Varicella					Varicella
Hepatitis A							HepA (2 doses)				HepA Series	
Meningococcal											MCV	
▲ Vaccine	Age ▶	Birth	1 Month	2 Months	4 Months	6 Months	12 Months	15 Months	18 Months	19-23 Months	2 - 3 Years	4 - 6 Years

## Reading the schedule

 This is the age or the age range the vaccine should be given. If a dose is missed, it should be given as soon as possible.

 These vaccines are for certain high-risk groups. Ask your doctor or nurse for more information.

 This dose is optional depending on the type of vaccine used. Ask your doctor or nurse for more information.

 The 4th dose of DTaP may be given as early as age 12 months. Ask your doctor or nurse for more information.

Adapted from the 2010 schedule of the U.S. Centers for Disease Control and Prevention, the American Academy of Pediatrics and the American Academy of Family Physicians. To find the recommended immunization schedule for ages 7 to 18 years, as well as information about adult immunizations, visit [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines)